

CHARACTERISTIC OF THE AMOUNT OF SOLAR ENERGY REACHING THE EARTH'S SURFACE FOR DIFFERENT PERIODS OF TIME IN THE STARA ZAGORA REGION.

Veselin Tashev¹ , Rolf Werner¹, Angel Manev¹, Mariana Goranova²

1. Space Research and Technology Institute (SRTI), Bulgarian Academy of Sciences, Stara Zagora Department

2. Technical University of Sofia, Faculty of Computer Systems and Management.

Abstract

The main source of solar energy falling on the Earth is the solar radiation. To measure the amount of solar energy use the data obtained from sensors solar radiation of the meteorological station Vantage Pro 2 Plus. The collected data from the sensors are integrated and recalculated in order to be obtained results for the solar energy that is absorbed for a certain period of time per unit Earth's surface. The purpose of research is to trace how great is the repeatability during different periods. A high annual repetition provides a good opportunity to forecast energy yields in the coming years. The monitoring had been carried out for the region of Stara Zagora.

Introduction

In the period of 5 years data for solar energy are measured and collected using the weather station Vantage Pro 2 Plus. During the 5 year time frame the data was captured every 15 minutes for total of more than 170 000 data points. The raw data was recorded by the solar radiation sensors of the station which was after that converted to solar energy absorbed over a given period of time per unit area. Quantitative accumulation of solar energy has been processed and aggregated in daily, monthly and annually periods. The results obtained are shown in graphic form. From the graphs shown the degree of repeatability during different periods.

Investigation of solar energy falling on the Earth.

As is well-known the solar radiation depends on many factors. Given that we can say that the solar energy which is directly related to the solar radiation would be different for the different places on the earth surface. This is based on the fact that only part of the solar energy is able to reach the earth surface. Some of the solar radiation is reflected right back into the space, some is absorbed by the stratosphere and troposphere. Overall, around 19% of the solar radiation does not reach the lower layers of the atmosphere (Roedel. 1994) [4].




Fig. 1 Solar energy measured for each day.

The The daily measurements of solar energy for 15 days in July for 5 years is shown in fig. 1. The theoretical values of the solar energy dependent on the direct solar radiation are marked by red bars and did not varied a lot, The measured solar energy, however, significantly varies in the different days as it is seen from the graph. For example, the values are almost the same for the 2nd and 7th days, while for the 1st and 3rd days the values are changed from 2000 to 7500 wh/m². Although the sun position is not significantly changed, the measured values differ due to actual atmosphere composition in the different days.

The atmosphere permeability is significantly affected by clouds, moisture, dust and other contaminants that absorb the solar energy.




Fig. 2 The average measured solar energy for every day for July.

In fig. 2, the daily measured average values of the solar energy in July are shown for 5 years. The daily deviation from the average value is marked by red lines. The dynamic change of the values of the 2nd and 7th days in comparison with the first and third day were significantly greater. Here even more clearly shows the changed dynamics due to changes in the atmosphere.




Fig. 3 Solar energy measured for each month from 2011 to 2016.

In fig.3, the solar energy values are measured and presented for the different months in a period of 5 years. The seasonal nature in the solar energy change is well followed. The solar energy is maximum in June and July and a minimum in January and December. It is worth to indicate that the dynamic change of the values for the corresponding months during the years is much less than the dynamic change of the values for the corresponding days as shown in fig 2.

Average deviation of the monthly solar energy




Fig. 4 Average measured solar energy for each month.

In fig. 4, the measured average solar energy per month in a period of 5 years is presented. The deviation from the average value for each month is given with red lines. It is well seen the smaller monthly deviation in comparison with the daily one. The figure shows that the smaller the deviations are during on the winter month January and during the summer month of August. This is due to the fact that these seasons are with more stable dynamic of the atmosphere.

Measured solar energy for each month




Fig . 5 Moving average with a period of 12 months for three years.

In fig. 5, a centered moving average value of solar energy for 12 months is presented for a period of about 5 years. It is shown in the chart with a red line. Monthly values of solar energy are shown with orange bars. As it is seen the value of moving average is substantially constant, i.e. the annual deviation is less than monthly one.

Conclusion

The research presented so far proved that the solar energy deviations for the corresponding months are less than the daily deviations, while the annual solar energy deviations are less than the monthly. Consequently, the calculated deviation of the corresponding values is smaller at a longer period of monitoring. From the obtained results we could state that it could be calculated and forecasted with a relative accuracy the expected energy from photovoltaic transformer for an annual period.

References

1. D a v I s I n s t r u m e n t s C o r p. Vantage Pro2 Plus. Referen. Guide.
2. J. M e e u s, Astronomische Algorithmen, Johann Ambrosius BarthVerlag, Leipzig-Berlin-Heidelberg, 1993
3. L e a n, J.L., Solar irradiance and climate forcing in the near future, Geophzs. Res. Lett. 28, pp. 4119-4122, 2001.
4. R o e d e l, W. Physik unserer Umwelt, 2. Aufl., Springer-Verlag Berlin Heidelberg, S.48, 1994.

**Contact address: engineer-physicist Veselin Tashev,
e-mail veselinlt@abv.bg**

Space Research and Technology Institute (SRTI) – BAS
Stara Zagora Department, Bulgaria.